


UNC School of Social Work  
Clinical Lecture Series

## Problem Solving Depression

January 23, 2017

Diane R. Dolan-Soto, MSW, LCSW  
Program Manager, UNC Internal Medicine  
Counseling Program  
Ambulatory Care Center, Chapel Hill

Diane.Dolan-Soto@unchealth.unc.edu 

---

---

---

---

---

---

---

---

## Agenda

- Information on Depression and Reasons for Treatment
- Evidenced Based Model for Depression Care
- Case Examples of Getting Better Through Problem Solving
- Build Knowledge and Skills to Fight Depression
- Steps to Move Through and Beyond Depression with Problem Solving
- Additional Tool: Unhelpful Thinking Styles and Depression
- Appendices:
  - Treatment Considerations
  - Behavioral Assessment Educational Aspects
  - Mnemonics for Depression

---

---

---

---

---

---

---

---

## Objectives

- Learn key aspects of a really successful approach to depression treatment.
- Learn steps to problem solving to reduce depression
- Come away from this training with information that you can use to incorporate in your practice to improve depression care and patient outcomes.

---

---

---

---

---

---

---

---

### Why Treat Depression in a Primary Care Setting?

- **Depression is a medical illness**
- 5-10% patients have MDDs. 7-16% lifetime risk in US adults
- 10% - 20% of adults visit their doctor during a depressive or anxious episode
  - Over 50% suffer from comorbid depressive or anxiety disorder
  - Comorbidity substantially increases medical utilization, is associated with greater chronicity, slower recovery and greater psychosocial disability
- 4<sup>th</sup> leading contributor to disability, expected to be 2<sup>nd</sup> by 2020
- People with diabetes, coronary artery disease, stroke, obesity, and HIV have a two-fold risk of depression compared to the general population

---

---

---

---

---

---

---

---

### Why Treat Depression in a Primary Care Setting?

- **One episode of depression increases likelihood of future episodes**
- **Patients *can* have increased pain related to depression**
  - Depression increases emotional *and* physical sensitivity

---

---

---

---

---

---

---

---

### Treatment Works!

- **80% of those who get treatment improve!**
- Only 50% of people with depression seek treatment
- **Effective treatment reduces symptoms and improves quality of life**
- 46% of adults improve with medication, 48% improve with counseling

**Providing *both* doubles the effectiveness of care**

---

---

---

---

---

---

---

---

### Evidenced Based Depression Care

- *Structured and integrated* care in a primary care setting
  - Uses trained physicians and counselors
  - Uses algorithms to *guide* treatment
  - Regular monitoring to assess and facilitate treatment progress
  - Builds on existing patient trust and access with provider/clinic
  - Example: IMPACT – Randomized Controlled Trial
    - For information on IMPACT: <http://aims.uw.edu/impact-improving-mood-promoting-access-collaborative-treatment>

---

---

---

---

---

---

---

---

### Evidenced Based Depression Care

- Past Usual Care
  - Seen by a doctor, usually not trained in depression care
  - See a counselor in another setting for long term talk therapy
  - May or may not see a psychiatrist for medication

---

---

---

---

---

---

---

---

### IMPACT Treatment and *Results!*

<b>Study year:</b>	
Doubled the effectiveness of usual care	JAMA 2002; 288: 2836-2845
Patients experienced better physical function	Callahan et al., JAGS 2005
Care was shown to benefit diverse populations	Arean et al. Medical Care 2005
<b>Beyond study:</b>	
Effects persisted one year after IMPACT	Hunkeler, et al., 2004 unpublished data
More depression-free days; 372 for IMPACT approach vs 265 for usual care patients	Hunkeler et al, under review (DFD calculation adapted from Lave et al., 1998)
As depression decreased for 1,001 arthritis patients, so did pain and pain interference	Lin et al, JAMA 2003

---

---

---

---

---


---

---

---

**PHQ-9 to Identify Depression**  
Patient Health Questionnaire-9

- Validated screener
  - Sensitivity = 91% for MDD,
  - Specificity = 89% for MDD
- 9 questions = 9 symptoms of depression
- 5 or more symptoms for 2 weeks could be depression
- Symptoms for 2+ months could be major depression
  - Rule outs
    - Medical conditions, medicines
    - Other issues that could look like depression: grief, illness/pain, some situational stressors, anxiety, etc.



*"Closest thing we've got to taking temperature for mood"*

---

---

---

---

---

---

---

---

---

---

---

---

PHQ-9 scoring, treatment (Tx) and follow up

#	Assessment and treatment		
<9	No or mild depression	No Tx	Screen in 1 yr or as needed
10-14	Moderate depression	Treatment Needed	RTC in 12 wks Education, Recommend Problem Solving, Activation, Rx?, Watchful monitoring
≥15	Severe depression	Treatment Needed	RTC in 4 wks Recommend Rx! and Problem Solving, Activation, Regular Monitoring

Clinically significant improvement = drop of ≥ 5 points  
Remission of depression is a drop of PHQ by 50%  
No depression ≤ 5 points

---

---

---

---

---

---

---

---

---

---

---

---

**Problem Solving Treatment\***

Behavioral health aspect of IMPACT:

**Three components of fighting depression**

- Focus on the Present
- Pleasant Activity
- Physical Activity

Teaches a structured approach to dealing with issues

Increases the ability to define problems and set realistic goals

Increases understanding of importance of **link between effort and mood**

\*From Dr. Jürgen Untzner's IMPACT Model

---

---

---

---

---

---

---

---

---

---

---

---

# Getting Better

## Case Examples of Problem Solving Depression

---

---

---

---

---

---

---

---

**S\*** Before

Referral: A 53 year old man with heart disease. Severely depressed related to health issues and limited mobility post stroke. Felt like he coped well with stroke, but felt like a burden to his family. Doctor thought he'd benefit from activating antidepressant such as bupropion, however Mr. S preferred to try counseling and increasing his activity first.

At Assessment: Strained relationship with his wife and family. Doing very little for himself. Most of his needs met through care from others. He was about to face the 'donut hole' with Medicare so he could only come for 3 to 4 visits including assessment.

Not patient's real initial. Elements of example have been changed to protect confidentiality.

- PHQ9 = 15
- Heart disease, other medical issues, recent stroke
- Family help with nearly all aspects of care
- Alienating supports
- Spending time alone: watching TV, crying or sleeping

---

---

---

---

---

---

---

---

**PROBLEM-SOLVING WORKSHEET**

Name: \_\_\_\_\_ Date: \_\_\_\_\_

Review of progress during previous week: \_\_\_\_\_

How have I handled you best with your effort? (0 - 10) (0 = Not at all, 10 = Extremely) \_\_\_\_\_

3. Problem: *Being dependent on others. Financial Autonomy. Support. Relationship do*

1. Goal: *Trying steps to be independent within reason*

3. Solutions:

Solution	Pros (+)	Cons (-)	Cost (Effort, Time, Money, Emotional Impact, Involving Others)
1) <i>Goal = better at similar as his own</i>	1) <i>Pros (+) What makes this a good choice? I'd feel like I wasn't a burden</i>	2) <i>Cons (-)</i>	A: 1.00 B: 1.00 C: 1.00 D: 1.00
2) <i>Help to get things going (eg. food bottles)</i>	3) <i>Pros (+) What makes this a good choice? Help me to be more independent</i>	3) <i>Cons (-)</i>	A: 1.00 B: 1.00 C: 1.00 D: 1.00
3) <i>Get on track by himself</i>	4) <i>Pros (+) What makes this a good choice? should...</i>	4) <i>Cons (-)</i>	A: 1.00 B: 1.00 C: 1.00 D: 1.00
4) <i>Should be on my own</i>	4) <i>Pros (+) What makes this a good choice? should...</i>	4) <i>Cons (-)</i>	A: 1.00 B: 1.00 C: 1.00 D: 1.00
5) <i>Get things done</i>	4) <i>Pros (+) What makes this a good choice? should...</i>	4) <i>Cons (-)</i>	A: 1.00 B: 1.00 C: 1.00 D: 1.00

---

---

---

---

---

---

---

---

**S\***

<p>Before</p> <ul style="list-style-type: none"> <li>• PHQ9 = 15</li> <li>• Heart disease, other medical issues, recent stroke</li> <li>• Family help with nearly all aspects of care</li> <li>• Alienating supports</li> <li>• Spending time alone: watching TV, crying or sleeping</li> </ul>	<p>After</p> <ul style="list-style-type: none"> <li>• PHQ9 = 7 (3 sess + phone 1x)</li> <li>• Engaging in self-care efforts</li> <li>• Improving relationships; reconnected to social and faith support network</li> <li>• Discovery of new measures of self-worth</li> </ul>
---	---

Not patient's real initial. Elements of example have been changed to protect confidentiality.

---

---

---

---

---

---

---

---

---

---

---

---

**K\***

Before

Referral: A 43 year old woman. Uncontrolled Type II Diabetes. Care for developmentally disabled 18 year old son and a 16 year old son.

At Assessment: Making efforts to take care of sons. Constant contact and requests for help from her mother. Little time with husband causing arguments. No time spent on self-care. Feels guilty about taking time for herself.

- PHQ9 = 18
- Uncontrolled diabetes, not taking Rx's regularly. Not physically active. Not sleeping well, over eating.
- Focus on others to exclusion of self-care

Not patient's real initial. Elements of example have been changed to protect confidentiality.

---

---

---

---

---

---

---

---

---

---

---

---

**PROBLEM-SOLVING WORKSHEET**

Name: \_\_\_\_\_ Date: 1/23/17

Review of progress during previous visit:

How Satisfied you feel with your effort (0 = Not at all, 10 = Extremely) \_\_\_\_\_ (Hand 0-10)

1. Problem: Not wanting more medical treatment Parental duties Insufficient resources

2. Task: Obtain self care with care for son Traveling Medical solutions

3. Solutions		4. Pros and Cons (Effort, Time, Money, Emotional Impact, Anxiety) (Others)				
		1) Cost (-)	2) Effort	3) Money	4) Anxiety	5) Other
1) <u>Go to bed earlier and not take calls</u>	1) <u>Pros (+) What makes this a good choice? <u>Don't sleep at all</u></u>	1) Cost (-)	2) Effort	3) Money	4) Anxiety	5) Other
2) <u>Go to bed earlier and not take calls</u>	2) <u>Pros (+) What makes this a good choice? <u>Have a clearer mind</u></u>	1) Cost (-)	2) Effort	3) Money	4) Anxiety	5) Other
3) <u>Write call on not to check messages</u>	3) <u>Pros (+) What makes this a good choice? <u>Don't feel so overwhelmed</u></u>	1) Cost (-)	2) Effort	3) Money	4) Anxiety	5) Other
4) <u>Get help to not message</u>	4) <u>Pros (+) What makes this a good choice? <u>Get help with calls</u></u>	1) Cost (-)	2) Effort	3) Money	4) Anxiety	5) Other
5) <u>Get help to not message</u>	5) <u>Pros (+) What makes this a good choice? <u>Don't feel so overwhelmed</u></u>	1) Cost (-)	2) Effort	3) Money	4) Anxiety	5) Other

---

---

---

---

---

---

---

---

---

---

---

---

Before	<b>K*</b>	After
<ul style="list-style-type: none"> <li>• PHQ9 = 18</li> <li>• Uncontrolled diabetes, not taking Rx's regularly. Not physically active. Not sleeping well, over eating.</li> <li>• Focus on others to exclusion of self-care</li> </ul>	<p><b>K*</b></p>	<ul style="list-style-type: none"> <li>• PHQ9 = 5 (at 11 mo, sess 12 of 12)</li> <li>• Taking Rx's regularly --Established healthy sleep hygiene and bedtime routine --Use of Rx's, PT, Mind Body Skills and regular walking. Better food portions.</li> <li>• Set healthy boundaries. Improved relationships. Getting out, socializing.</li> <li>• Surprise benefit re: diabetes</li> </ul>

Not patient's real initial. Elements of example have been changed to protect confidentiality.

---

---

---

---

---

---

---

---

## Behavioral Approach: Build Knowledge and Skills to Fight Depression

Sharing What I've Learned for Effective Depression Care

---

---

---

---

---

---

---

---

<p style="text-align: center;"><b>Overview of Behavioral Depression Care</b></p> <p style="text-align: center;"><b>Assessment Visit</b></p> <p>1<sup>st</sup> visit: assessment, diagnosis, education as appropriate, and treatment planning. Format to the right:</p>	<ul style="list-style-type: none"> <li>• <b>Assess</b> <ul style="list-style-type: none"> <li>• Full psychosocial diagnostic</li> <li>• Risk</li> <li>• PHQ9</li> <li>• Mood symptoms <i>emphasis on functionality and efforts</i></li> </ul> </li> <li>• <b>And Educate!</b> <ul style="list-style-type: none"> <li>• On Depression and PHQ9</li> <li>• Treatment options</li> <li>• Link impact of effort to mood</li> </ul> </li> <li>• <b>Introduce Behavioral Approach</b> <ul style="list-style-type: none"> <li>• 3 Components to fight depression</li> <li>• Self-report measure - <i>"If we're doing the right things the PHQ9 score will be going down. If it's not, it means you and I need to change the goals, you and your doctor need to make changes, or we made need to involve a psychiatrist or other supports or treatments."</i></li> </ul> </li> <li>• <b>Set Goal for Treatment and Return</b> Appointment for First Counseling Visit</li> </ul>
--	---

---

---

---

---

---

---

---

---

## Share the Diagnosis!

- A diagnosis is based on symptoms. Helpful for billing and a focus on what we're going to work on.
- "A diagnosis is more like a street sign, it's where you are, *not* who you are." – Diane R. Dolan-Soto, LCSW
- Reduce stigma
- Build trust, rapport and engagement, *and understanding of what they need to be working on*

---

---

---

---

---

---

---

---

## "Depression is like Pneumonia"

### Pneumonia

- 1 episode makes you more susceptible to getting it in the future.
- Medication for about a month
- Work with your doctor and make efforts to get better
- Can take 1 to 4 months to get better

### Depression

- 1 episode makes you more susceptible to getting it in the future.
- May take both medication, like with severe depression, and effort to get better.
- Life cycle is 9 to 12 months

---

---

---

---


---

---

---

---

## Depression = Downward Spiral




---

---

---

---


---

---

---

---






---

---

---

---

---

---

---

---

- ## Steps to Move Through and Beyond Depression
- Orient to Treatment
  - Keys for Moving Forward
  - How To Define The Problem and Set a Goal
  - How to Figure Out Solutions
  - Evaluate Pros and Cons
  - How To Do More Successful Goal Setting
  - Move Beyond Depression and Prevent Relapse

---

---

---

---

---

---

---

---

### Overview of Behavioral Depression Care

#### Return Visit

Return sessions follow format to the right:

- **First Visit: Orient to Treatment. Ongoing:**
- **Check-In**
  - PHQ9
  - Rate mood on scale of 0-10
  - Rate satisfaction with effort 0-10
- **Review / Assess**
  - Risk
  - **Functional changes:** medication sleeping, eating, substance use, etc.
  - **Efforts, progress, challenges** related to previous visit's goals
- **Aid treatment change, linkage, etc.**
- **Link impact of effort to mood**
- **Problem Solving Focus**
  - Presenting issue or problem patient wants to address
- **Set Goals and Return Timeframe**
  - Identify Solutions – set manageable goals
  - Pleasant and Physical Activity goals
- **Continue to Educate and Encourage!!**

---

---

---

---

---

---

---

---

## Keys to Move Forward

1. Focus on the present *and* what's within your control

**“What's the *smallest* thing you can do to feel good for the effort?”**

— Joseph B. Quinn, MSW

1. Pleasant Activity – At least 2x week

**Feel better *while* doing, “Not a should”**

2. Physical activity – 3x week ‘as doable’

For providers: Note emphasis on CBT and Solution Focused aspects

### Goal Setting Initial Aims

= Identify easiest goal(s) that will result in a successful outcome

**BUILD HOPE *and* CONFIDENCE**

---

---

---

---

---

---

---

---

---

---

**PROBLEM-SOLVING WORKSHEET**

Name: \_\_\_\_\_ Date: \_\_\_\_\_ Title: \_\_\_\_\_

Review of progress during previous week: \_\_\_\_\_

Rate how satisfied you feel with your effort (1 = Not at all, 10 = Extremely): \_\_\_\_\_ Mood (1-10): \_\_\_\_\_

1. Problem: \_\_\_\_\_ Perceived (1-10): \_\_\_\_\_

2. Goal: \_\_\_\_\_

3. Solutions:

4. Pros versus Cons (Effort, Time, Money, Emotional Impact, Involving Others)

	a) Pros (+)	What makes this a good choice?	b) Cons (-)	Effort	Sat.	Medium	Act
1)							
2)							
3)							
4)							
5)							

---

---

---

---

---

---

---

---

---

---

## Get a Baseline

- Rate Mood
  - 1 to 10 with 10 being the best – Initially “Don’t make it pretty!”
  - The more realistic the rating, the more your person will trust when the numbers improve
  - Quantify how your person is feeling now
- Rate Satisfaction with Effort in taking care of mood
  - 1 to 10 with 10 being the best
  - Quantify how satisfied your person is and how comfortable/uncomfortable they are with current situation; also identifies motivation for change

---

---

---

---

---

---

---

---

---

---

## How To Define The Problem

“Depressed thinking is like having blinders on.” –only see the negative, everything looks too difficult.

- Identify the problem *of the moment!*
  - ‘My home is a disaster.’
- Set an *overarching* goal – What do you want to happen?
  - ‘I want to reclaim my home.’

---

---

---

---

---

---

---

---

**PROBLEM-SOLVING WORKSHEET**

Name: \_\_\_\_\_ Date: \_\_\_\_\_ Title: \_\_\_\_\_

Review of progress during previous week: \_\_\_\_\_

Rate how satisfied you feel with your effort: 10 (Not at all) to 100 (Extremely) \_\_\_\_\_ Mood 10-100 \_\_\_\_\_

1. Problem: Mail and other papers piling up (Present/As-to-Be?) \_\_\_\_\_

2. Goal: Reduce backlog and create some order

3. Solutions:

4. Pros versus Cons (If Pros, Time, Money, Emotional Impact, Involving Others)	5. Chosen Solution
(a) Cons (+)	Pros Cons Time Money Emotional Impact Involving Others
(b) Cons (-)	Pros Cons Time Money Emotional Impact Involving Others
(c) Cons (+)	Pros Cons Time Money Emotional Impact Involving Others
(d) Cons (-)	Pros Cons Time Money Emotional Impact Involving Others

---

---

---

---

---

---

---

---

## How to Figure Out Solutions

- Look for possible solutions toward **one aspect** of the goal.
- Think small!!
  - What's within your control?
  - Where would you start?
  - Who or what would be needed?
  - What could you do in 5 to 20 minutes?

---

---

---

---

---

---

---

---

## How to Figure Out Solutions

- Identify at least 2 to 5 possible solutions
  - It helps to recognize there are options to “open the blinders”
  - This may sound simplistic. For your patient, it’s not!
  - You may need to reassure your client/patient: “If this was easy you would’ve already figured it out.”

---

---

---

---

---

---

---

---

**PROBLEM-SOLVING WORKSHEET**

Name: \_\_\_\_\_ Date: \_\_\_\_\_ Title: \_\_\_\_\_

Review of progress during previous week: \_\_\_\_\_

State how satisfied you feel with your effort: 1=Not at all, 10=Extremely. \_\_\_\_\_ Mood: 1=Bad, 10=Good \_\_\_\_\_

1. Problem: Mail and other papers piling up "Desired Action?" \_\_\_\_\_

2. Goal: Reduce backlog and create some order

3. Solution:	4. Pros/Cons (Rate, Time, Money, Emotional Impact, Involving Others)	5. Rating (1-5)
Sort papers into piles (maybe w/ a friend)		
Pull out and shred junk mail!		
Go thru and pull out any bills		
Put file for bills, shredder & recycling bag near door & sort when it comes in		

---

---

---

---

---

---

---

---

## Identify the Pros

- Repeat back each solution and ask:
  - “What makes this a good choice for you?”
- Helps literally with structuring their thought process
- Look for “me” and “I” based answers. “It would make me feel good...” “I like...” “I want...”
  - These are usually the most powerful motivators.
- Often ‘shoulds’ are the first response
  - Try to avoid shoulds. More likely to be difficult. May compound feelings of hopelessness. Less likely to motivate.

---

---

---

---

---

---

---

---

**PROBLEM-SOLVING WORKSHEET**

Name: \_\_\_\_\_ Date: \_\_\_\_\_ TIME: \_\_\_\_\_

Review of progress during previous week:

Rate how satisfied you feel with your effort (0 = Not at all, 10 = Extremely): \_\_\_\_\_ Mood (0-10): \_\_\_\_\_

1. Problem: **Mail and other papers piling up** Present Intensity? \_\_\_\_\_

2. Goal: **Reduce backlog and create some order**

3. Solution:

3. Solution:	4. Pros versus Cons (Effort, Time, Money, Emotional Impact, Involving Others)												
a) Sort papers into piles (maybe w/ a friend)	<table border="1"> <tr> <td>4) Pros (+)</td> <td>5) What makes this a good choice?</td> <td>6) Cons (-)</td> <td>A Little</td> <td>Medium</td> <td>A Lot</td> </tr> <tr> <td></td> <td>I'd feel more organized</td> <td></td> <td></td> <td></td> <td></td> </tr> </table>	4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot		I'd feel more organized				
4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot								
	I'd feel more organized												
b) Pull out and shred junk mail!	<table border="1"> <tr> <td>4) Pros (+)</td> <td>5) What makes this a good choice?</td> <td>6) Cons (-)</td> <td>A Little</td> <td>Medium</td> <td>A Lot</td> </tr> <tr> <td></td> <td>I need to do this</td> <td></td> <td></td> <td></td> <td></td> </tr> </table>	4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot		I need to do this				
4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot								
	I need to do this												
c) Go thru and pull out any bills	<table border="1"> <tr> <td>4) Pros (+)</td> <td>5) What makes this a good choice?</td> <td>6) Cons (-)</td> <td>A Little</td> <td>Medium</td> <td>A Lot</td> </tr> <tr> <td></td> <td>I'd stop worrying that I missed something. I'd feel relieved.</td> <td></td> <td></td> <td></td> <td></td> </tr> </table>	4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot		I'd stop worrying that I missed something. I'd feel relieved.				
4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot								
	I'd stop worrying that I missed something. I'd feel relieved.												
d) Put file for bills, shredder & recycling bag near door & sort when it comes in	<table border="1"> <tr> <td>4) Pros (+)</td> <td>5) What makes this a good choice?</td> <td>6) Cons (-)</td> <td>A Little</td> <td>Medium</td> <td>A Lot</td> </tr> <tr> <td></td> <td>It would make things easier, prevent future pile up of papers.</td> <td></td> <td></td> <td></td> <td></td> </tr> </table>	4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot		It would make things easier, prevent future pile up of papers.				
4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot								
	It would make things easier, prevent future pile up of papers.												

---

---

---

---

---

---

---

---

---

---

---

---

## Identify the Cons

State the solution, the 'pro' and then help your client evaluate the cons for each solution.

Cons (-)	A Little	Medium	A Lot
Effort	[ ]	[ ]	[ ]
Time	[ ]	[ ]	[ ]
Money	[ ]	[ ]	[ ]
Emotional Impact	[ ]	[ ]	[ ]
Involving Others	[ ]	[ ]	[ ]

---

---

---

---

---

---

---

---

---

---

---

---

**PROBLEM-SOLVING WORKSHEET**

Name: \_\_\_\_\_ Date: \_\_\_\_\_ TIME: \_\_\_\_\_

Review of progress during previous week:

Rate how satisfied you feel with your effort (0 = Not at all, 10 = Extremely): \_\_\_\_\_ Mood (0-10): \_\_\_\_\_

1. Problem: **Mail and other papers piling up** Present Intensity? \_\_\_\_\_

2. Goal: **Reduce backlog and create some order**

3. Solution:

3. Solution:	4. Pros versus Cons (Effort, Time, Money, Emotional Impact, Involving Others)												
a) Sort papers into piles (maybe w/ a friend)	<table border="1"> <tr> <td>4) Pros (+)</td> <td>5) What makes this a good choice?</td> <td>6) Cons (-)</td> <td>A Little</td> <td>Medium</td> <td>A Lot</td> </tr> <tr> <td></td> <td>I'd feel more organized</td> <td></td> <td></td> <td></td> <td></td> </tr> </table>	4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot		I'd feel more organized				
4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot								
	I'd feel more organized												
b) Pull out and shred junk mail!	<table border="1"> <tr> <td>4) Pros (+)</td> <td>5) What makes this a good choice?</td> <td>6) Cons (-)</td> <td>A Little</td> <td>Medium</td> <td>A Lot</td> </tr> <tr> <td></td> <td>I need to do this</td> <td></td> <td></td> <td></td> <td></td> </tr> </table>	4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot		I need to do this				
4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot								
	I need to do this												
c) Go thru and pull out any bills	<table border="1"> <tr> <td>4) Pros (+)</td> <td>5) What makes this a good choice?</td> <td>6) Cons (-)</td> <td>A Little</td> <td>Medium</td> <td>A Lot</td> </tr> <tr> <td></td> <td>I'd stop worrying that I missed something. I'd feel relieved.</td> <td></td> <td></td> <td></td> <td></td> </tr> </table>	4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot		I'd stop worrying that I missed something. I'd feel relieved.				
4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot								
	I'd stop worrying that I missed something. I'd feel relieved.												
d) Put file for bills, shredder & recycling bag near door & sort when it comes in	<table border="1"> <tr> <td>4) Pros (+)</td> <td>5) What makes this a good choice?</td> <td>6) Cons (-)</td> <td>A Little</td> <td>Medium</td> <td>A Lot</td> </tr> <tr> <td></td> <td>It would make things easier, prevent future pile up of papers.</td> <td></td> <td></td> <td></td> <td></td> </tr> </table>	4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot		It would make things easier, prevent future pile up of papers.				
4) Pros (+)	5) What makes this a good choice?	6) Cons (-)	A Little	Medium	A Lot								
	It would make things easier, prevent future pile up of papers.												

---

---

---

---

---

---

---

---

---

---

---

---

## Physical and Pleasant Activity

- Physical Activity
  - **Make the goal concrete:** eg. walk 3 times a week for 15-20 minutes
  - Be creative! It all counts!
- Pleasant Activity
  - Feel good while you're doing, lose track of time. Music, art, crafts, games, social, etc.
  - Passive activities can support background worrying, don't usually improve mood and functioning.
    - Too much TV/computer can contribute to depression -- news, crime shows, negative comparison to what others are doing/have
- If they don't have what they need *but they want to get it*, that may be the first goal.
  - Every step takes effort, a precious commodity with depression

---

---

---

---

---

---

---

---

## How To Do More Successful Goal Setting

### Assess ability to actually accomplish the goal!

ASK: "On a scale of 1-10 how likely are you to be able to do \_\_\_ with 10 being the most likely?"

Rated 1-7 = **Rework the goal**

"What would it take to make it an 8?"

Rated 8 or above = **most likely success!**

Make an appointment!

Increases the likelihood for a successful goal.

What's the best day and time - What else will help make the goal possible

---

---

---

---

---

---

---

---

## Goal Sheet

### Goals until next visit:

Today we discussed your goal of: "Reduce backlog of papers and create some order"

Today you set step(s) below for yourself to work towards this goal.

Take 20 minutes Saturday after lunch to go through and pull out any bills, consider listening to favorite upbeat CD

Pleasant Activity: Call my friend and / or go to lunch together

---

---

---

---

---

---

---

---

## Goal Sheet

Physical Activity: Walk at home or at work during lunch for 10-15 minutes 2-3 times a week.

**Problem Solving weekly goal reminder:**

1. Focus on the present and what's within your control  
Tasks and problems - pick the smallest thing you can do that will make you feel good for the effort. Focus only on what's within your control
2. Pleasant activity 1-2 times a week
3. Physical activity 3 times a week, within what's physically comfortable

---

---

---

---

---

---

---

---

## To Keep It Moving Forward

Steps to Move Through and Beyond Depression

- **Review:** "Tell me what you've been doing to take care of your mood."
- "Focus on what someone does, not what they haven't done"
- If you set goals, **review them!!** Link impact of effort to mood
- Continue to help clients/patients to:
  - Monitor/rate. Share PHQ and treatment progress
  - With realistic expectations, judgments
  - Learn to pace their efforts
  - Recognize and sustain their progress
  - Discuss what relapse would look like and discuss how to prevent relapse
  - Determine next steps *with* your client

---

---

---

---

---

---

---

---

## Unhelpful Thinking Styles

Adding a Powerful Tool to Your Toolkit

---

---

---

---

---

---

---

---

**“Anything look familiar?”**

Identify which unhelpful thinking styles are used

---

---

---

---

---

---

---

---

**Work on Unhelpful Thinking Styles**

- Initial thought – Immediate real concern or risk? Address!!
- If no immediate risk, then assess for unhelpful thinking
  - Evidence for thought? Evidence against thought?
- For an unhelpful thought, develop a new, revised, more realistic and balanced thought based on the evidence
- Practice work with unhelpful thinking styles
  - Notice. *Kindly* interrupt. [Over time learn to] Change.
    - Repeat effort. Change comes with practice.
    - Work to get rid of the ANTs! (Automatic Negative Thoughts)

---

---

---

---

---

---

---

---

**Problem Solving Depression**

Adding Powerful Skills to Successfully Fight Depression

Diane.Dolan-Soto@unchealth.unc.edu

---

---

---

---

---

---

---

---



## Acknowledgements

- **IMPACT**
  - Improving Mood Promoting Access to Collaborative treatment – evidenced based depression treatment for primary care. Problem Solving Treatment. Dr. Jürgen Unützer. <http://aims.uw.edu/impact-improving-mood-promoting-access-collaborative-treatment>
- **UNC General Internal Medicine, Ambulatory Care Center**
  - Robert Malone, PharmD
  - Dr. Michael Pignone
  - Dr. Amy Weil
  - Betsy Shilliday, PharmD
  - Dr. Shana Ratner
  - Internal Medicine Psychiatrist Consultant Dr. Robin Reed
  - Madeline Waife
  - Deborah Barrett, PhD, LCSW
  - Laura Cruz

---

---

---

---

---

---

---

---

---

---

## References

### Primary References

- UNC Internal Medicine Preclinic Conference Training for residents: Depression: Screening, Diagnosis and Treatment. Amy Weil MD, Diane R. Dolan-Soto LCSW, Robin Reed MD. 2015
- In the Clinic: Depression. Annals of Internal Medicine. May 2007;146(9):IT51-IT516. Available from: Academic Search Premier, Ipswich, MA. Accessed November 4, 2009.
- Carlat, D. The Psychiatric Review of Systems: A Screening Tool for Family Physicians. AAFP 58:1627-1635 (1998).
- UNC Depression Care Program Updated Screening and Treatment Algorithms.

---

---

---

---

---

---

---

---

---

---

## References

### Additional References

- Carlat, Daniel J M.D. The Psychiatric Review of Symptoms: A Screening Tool for Family Physicians. American Family Physician. Nov. 1, 1998. Am Fam Phys Web.
- Diabetes and Depression, NIH Publication No.02-5003 from the National Institute of Mental Health. [http://www.dlife.com/diabetes/information/daily\\_living/depression\\_and\\_coping/](http://www.dlife.com/diabetes/information/daily_living/depression_and_coping/)
- Healthline. Diabetes, Alcohol, and Social Drinking. Found online 3/11/16. <http://www.com/health/type-2-diabetes/facts-diabetes-alcohol#1>

---

---

---

---

---

---

---

---

---

---

## References

- **Additional References**
- Kroenke K. A 75-year-old man with depression. *Journal of the American Medical Association*. 2002; 287: 1568-1576.
- O'Connor, E. A., Whitlock, E. P., Beil, T. L., & Gaynes, B. N. (2009). Screening for depression in adult patients in primary care settings: a systematic evidence review. *Annals of Internal Medicine*, 151(11), 793-803. doi: 10.1059/0003-4819-151-11-200912010-00007
- University of Michigan (UMHS) Depression Guideline Update, October 2005

---

---

---

---

---

---

---

---

## Appendix A Treatment Considerations

What to Consider *Before* Starting Treatment  
Medication  
Indications for Additional or Alternate Treatment

---

---

---

---

---

---

---

---

### *Before* Starting Treatment

- **Check for Rule Outs**
  - Check thyroid function
  - Consider other medical conditions / medicines / issues
- **Bipolar Disorder brief screen:**
  - Have you had periods of feeling so happy or energetic that your friends told you that you were talking too fast or that you were too 'hyper'?<sup>4</sup>
    - If yes – *And* how many nights have you been able to go without sleep *and* had plenty of energy?
    - If yes – *And* while being involved in risky behaviors?
- **Assess for and Address Risk**
  - Suicidality, substance abuse, dependence
  - Presence of violence, etc.

---

---

---

---

---

---

---

---

## Medication

- Antidepressants (ADs), SSRIs and SNRIs, can cause activation and agitation when first started, may temporarily exacerbate anxiety before mood and functioning improve.
  - No specific AD is more effective than another.
  - Consider cost, side effects, drug-drug interactions, and/or family/personal history w/ ADs
  - Start low and increase gradually to effective dose as soon as tolerable
  
- May help with moderate depression, **most effective for severe depression**
  - Within  $\pm$  6 weeks, half receiving ADs have at least 50% reduction in sx
  - Tx timeframe 9-12 months; can be up to 24 months, or ongoing depending on severity, history, personality organization...
  - ADs produce full remission in 30% of patients with MDD and partial response, a 50% reduction in PHQ-9 score, in 60%

---

---

---

---

---

---

---

---

---

---

## Indicators for Additional or Alternate Treatment

- Active risk – suicidality, other self-harm behaviors, homicidality
- Active substance abuse
- Seek psychiatric consult when:
  - Severe depression and no onsite option for pharmacotherapy
  - More than 2-3 medication trials without significant reduction in symptoms or improvement.
  - Multiple medications without benefit
  - Longstanding depression without benefit from prior treatment
  - Complicating, multiple diagnoses – for example: Depression and PTSD
- Linkage to outside CBT when not available at your setting

---

---

---

---

---

---

---

---

---

---

## Appendix B Assess and Educate

Psychoeducational Components of Behavioral  
Depression Education

---

---

---

---

---

---

---

---

---

---

## Assess *and* Educate

- **Healthy Self-Care Management is Essential!!**  
As appropriate, Teach and Share Information!
- Help your client learn how to positively affect mood and functioning
- Help your client identify realistic / unrealistic expectations and judgments

---

---

---

---

---

---

---

---

## Assess *and* Educate

- **Sleep** (examples)
  - Poor sleep worsens mood and functioning (and is a significant risk factor for suicidality)
  - Electronics adversely impact sleep – TV, phone, pad, computer.
    - Can be activating. Use of smartphone or pad can interrupt melatonin production for 1-2 hours.
  - Caffeine –half-life of 5-9 hrs; caffeine after 12 – 2 pm can affect sleep
  - Efficiency: # of hrs asleep ÷ # of hrs in bed x 100% if ≤ 75% = restlessness
  - Resource: SHUTi – Online CBT for insomnia program [www.myshuti.com](http://www.myshuti.com)

---

---

---

---

---

---

---

---

## Assess *and* Educate

- **Eating, nutrition** (examples)
  - Food provides the building blocks for our emotions and energy
  - Food benefits 'last' in the body for about 5 hours
  - **Breakfast, lunch and dinner?**
    - Eating regularly helps mood and functioning
    - Skipping meals may contribute to dips in mood, irritability, over eating, gaining or holding onto weight
    - Eating breakfast can help to regulate body weight
    - Eating late at night lowers appetite in the morning, may also affect sleep

---

---

---

---

---

---

---

---

### Assess *and* Educate

- Substance Use / Abuse / Dependence (examples)
  - Caffeine – can add to irritation and muscle tension in addition to affecting sleep
  - Cigarettes – can be emotionally soothing, but are physically activating
  - Alcohol – central nervous system depressant, can affect the body up to 3 days, sleep interruptor; can ‘out weigh’ benefit of ADs

---

---

---

---

---

---

---

---

### Assess *and* Educate

- Marijuana – helps to disconnect from mood; withdrawal response within 4 days, can cause anxiety, reinforce need/addiction
- ...and impact of other substances. Self-medication is ‘medication’ with unclear amounts, action and effects.
  - Can interfere with and / or reduce effectiveness of ADs.
  - Substance use/abuse/dependence may be a significant contributor to depression

---

---

---

---

---

---

---

---

### Assess *and* Educate

- Physical activity (examples)
  - It all counts!
    - Cleaning, shopping, swimming, dancing, running , playing with kids, critters.
  - Walking for 10 minutes raises mood and energy for 2 hours
  - Walking for 20 to 30 minutes, 3 to 5 times a week, is equivalent to an antidepressant
  - The way we think about what we do can have a large impact on the benefit we get from the physical activity we engage in.

---

---

---

---

---

---

---

---

### Additional: Depression MNEMONICS

Mania – DIGFAST	Depression – SIGECAPS ≥ 2 months	Dysthymia – He's 2 SAD ≥ 2 years
Distractability	Sleep disorder	Hopelessness
Indiscretion (Pleasurable Activities)	Interest Deficit (anhedonia)	Energy loss or fatigue
Grandiosity	Guilt	Self-esteem is low
Flight of Ideas	Energy Deficit	2 yr min depressed mood most times/days
Activity Increased	Concentration Deficit	Sleep increased/decreased
Sleep Decreased	Appetite Disorder	Appetite increased/decreased
Talkativeness (Pressured Speech)	Psychomotor Agitation/Retardation	Decision-making or concentration impaired
	Suicidality	

---



---



---



---



---



---



---



---